ELSEVIER

Volume 239, Numbers 1–2, November 1995

Inorganica Chimica Acta

Contents

Preliminary Communication

Homoleptic Fe(II) complexes of new 2,2′:6′,2″-terpyridine ligands with 4′-*p*-hydroquinonyl or 4′-*p*-quinonyl pendants 1
G.D. Storrier, S.B. Colbran, D.B. Hibbert (Sydney, NSW, Australia)

Articles

Magnetic and spectroscopic properties of a series of linear homo- and heterotrinuclear metal compounds with asymmetric 3,4-disubstituted 1,2,4-triazoles as ligands 5
P.J. van Koningsbruggen, J.G. Haasnoot, W. Vreugdenhil, J. Reedijk (Leiden, Netherlands), O. Kahn (Orsay, France)

Zinc complexes of dipyridyl-triazoles. 13
U. Hartmann, H. Vahrenkamp (Freiburg, Germany)

Synthesis and some properties of T-cage type $Co^{III}_4Hg^{II}_4$ and $Rh^{III}_4Hg^{II}_4$ octanuclear complexes with 2-aminoethanethiolate 19
Y. Kageyama, T. Konno, K. Okamoto (Ibaraki, Japan), J. Hidaka (Fukuoka, Japan)

Molecular complex design: bridging the tetrakiscarboxylatodirhodium core 25
D.F. Taber, R.P. Meagley, J.P. Louey, A.L. Rheingold (Newark, DE, USA)

Electrospray mass spectrometry and X-ray crystallography studies of divalent metal ion complexes of tris(2-pyridylmethyl)amine 29
C.S. Allen, C.-L. Chuang, M. Cornebise, J.W. Canary (New York, NY, USA)

Synthesis and characterization of the polyoxothioanion α-$[PW_{11}NbSO_{39}]^{4-}$ derived from the Keggin structure 39
E. Cadot, V. Béreau, F. Sécheresse (Versailles, France)

Lead(II) complexes of 1,4,7,10-tetraazacyclododecane-*N*,*N*′,*N*″,*N*‴-tetraacetate: solution chemistry and application to tumor localization with ^{203}Pb labeled monoclonal antibodies 43
C.G. Pippin, T.J. McMurry, M.W. Brechbiel, M. McDonald (Bethesda, MD, USA), R. Lambrecht (Riyadh, Saudi Arabia), D. Milenic, M. Roselli (Bethesda, MD, USA), D. Colcher (Bethesda, MD, SA and Omaha, NE, USA), O.A. Gansow (Bethesda, MD, USA)

New examples of dinuclear copper complexes with ferromagnetic interactions mediated by μ-1,1 bridging azido and nitrito groups: structures and magnetic properties of $[L_2(N_3)_2Cu_2]$ and $[L_2(NO_2)_2Cu_2]\cdot H_2O$ (L = 7-amino-4-methyl-5-aza-3-hepten-2-onato(1−)) 53
J.P. Costes, F. Dahan, J. Ruiz, J.P. Laurent (Toulouse, France)

Coordinatively saturated Pt(II) complexes as potential precursors of organometallic polymers 61
F. Giordano, B. Panunzi, A. Roviello, F. Ruffo (Naples, Italy)

A thermogravimetric, single crystal X-ray, and spectroscopic study of several iron(III) halide adducts with crown ethers 67
U. Russo, B. Zarli, G. Favero, G. Valle (Padua, Italy), G.J. Long (Rolla, MO, USA)

Crystal structures of *trans*-$[Ru(dppe)_2(CO)(Cl)](BF_4)\cdot 2(toluene)$ and *trans*-$[Ru(dppm)_2(CO)(Cl)](BF_4)\cdot CH_2Cl_2$: a study of the steric and electronic ligand effects of *trans*-positioned diphosphine ligands 77
L.F. Szczepura (Buffalo, NY, USA), J. Giambra (Fredonia, NY, USA), R.F. See (Buffalo, NY, USA), H. Lawson, T.S. Janik (Fredonia, NY, USA), A.J. Jircitano (Erie, PA, USA), M.R. Churchill, K.J. Takeuchi (Buffalo, NY, USA)

The reaction of cadmium dicyclohexyldithiophosphinate with tributylphosphine. The crystal and molecular structures of tetrakis(dicyclohexyldithiophosphinato)dicadmium(II) chloroform solvate, $[Cd_2(S_2PCy_2)_4]\cdot CHCl_3$, and of the tributylphosphine adduct $[Cd(S_2PCy_2)_2(PBu_3)]$ 87
J.S. Casas, M.S. García-Tasende, A. Sánchez, J. Sordo, E.M. Vázquez-López (Santiago de Compostela, Galicia, Spain), E.E. Castellano, J. Zukerman-Schpector (São Carlos, SP, Brazil)

Elsevier Science S.A.

Coordination chemistry of iminodiacetic esters. Structural and spectroscopic characterization of copper(II) and zinc(II) monoethyliminodiacetato complexes 93
J.A. Guevara-García, N. Barba-Behrens, A.R. Tapia-Benavides, M.J. Rosales-Hoz, R. Contreras (Mexico, D.F., Mexico)

Kinetics of the reaction of chromium(II) with ethyliodoacetate, iodoacetatopentaamminecobalt(III) 99
A.D. Jordan, S.L. Scott, R.B. Jordan (Edmonton, Alta., Canada)

Binuclear copper(II) complexes containing phenols and catechols 107
R.J. Butcher (Washington, DC, USA), G. Diven, G. Erickson (Wollongong, NSW, Australia), J. Jasinski (Keene, NH, USA), G.M. Mockler (Wollongong, NSW, Australia), R.Y. Pozdniakov (Keene, NH, USA), E. Sinn (Kingston-upon-Hull, UK)

Structure, electrochemical and dynamic behavior of bis(pentamethylcyclopentadienyl)titanium(IV)dithiolene complexes 117
F. Guyon (Besancon, France), M. Fourmigué (Nantes, France), P. Audebert, J. Amaudrut (Besancon, France)

Synthesis and characterisation of the six-membered cyclopalladated complexes of 2-benzylbenzothiazole 125
Y. Fuchita, K. Yoshinaga, H. Kusaba (Fukuoka, Japan), M. Mori, K. Hiraki (Nagasaki, Japan), K. Takehara (Fukuoka, Japan)

Cationic diamineplatinum(II) complexes of nalidixic acid 133
Sung Si Lee, Ok-Sang Jung (Seoul, South Korea), Chong Ock Lee, Sang Un Choi (Taejon, South Korea), Moo-Jin Jun, Youn Soo Sohn (Seoul, South Korea)

Spectroscopic study of ion binding in synthetic polyelectrolytes using lanthanide ions 139
S. Lis, Z. Wang, G.R. Choppin (Tallahassee, FL, USA)

Oxovanadium(IV) complexes of citric and tartaric acids in aqueous solution 145
T. Kiss, P. Buglyó (Debrecen, Hungary), D. Sanna, G. Micera (Sassari, Italy), P. Decock, D. Dewaele (Dunkerque, France)

Notes

Aliphatic cyclopalladation. Crystal structure of $[\overline{Pd(1\text{-}CH_2\text{-}2\text{-}\{HC{=}N\}[2',6'\text{-}(CH_3)_2C_6H_3]}\text{-}3,5\text{-}(CH_3)_2C_6H_2)Br]_2$ 155
G. De Munno, M. Ghedini, F. Neve (Arcavacata di Rende, CS, Italy)

Synthesis and crystal structure of a tetranuclear bis(diphenylphosphino)methane palladium complex capped by a μ_4-bonded sulfide: $[Pd_4(\mu\text{-dppm})_4(\mu_4\text{-S})]\cdot 4(CH_3)_2CO$ 159
M. Maekawa, M. Munakata, T. Kuroda-Sowa, T. Goto (Osaka, Japan)

Crystal structure and magnetic properties of manganese(II) mellitate, $[Mn_2\{C_6(COO)_6\}(H_2O)_6][Mn(H_2O)_6]\cdot 2H_2O$ with two-dimensional layered structure and three-dimensional hydrogen bonding networks 165
L.P. Wu, M. Yamamoto, T. Kuroda-Sowa, M. Maekawa, J. Fukui (Osaka, Japan), M. Munakata (Osaka, Japan and Okazaki, Japan)

Crystal structures of two *N*-methyltriphenylphosphoranimines: triphenylphosphine-*N*-methylimine borane, $Ph_3PN(CH_3)BH_3$ and *N,N*-dimethylaminotriphenylphosphonium tetrafluoroborate, $[Ph_3PN(CH_3)_2]^+BF_4^-$ 171
W.K. Holley, G.E. Ryschkewitsch, A.E. Koziol, G.J. Palenik (Gainesville, FL, USA)

Adducts of organotin(IV), tin(IV) and tin(II) halides with 1-methyl-imidazoline-2(3*H*)-thione (Hmimt) and imidazoline-2(1,3*H*)-thione (Himt). Synthesis, spectroscopic (IR, Mössbauer and ^1H, ^{13}C, ^{119}Sn NMR) studies and in vitro antitumour activity 177
D. Kovala-Demertzi, P. Tauridou (Ioannina, Greece), U. Russo (Padua, Italy), M. Gielen (Brussels, Belgium)

Stereoelectronic properties of the sulfinyl-substituted phosphine ligands: synthesis and X-ray structure of the stable complex $[PdCl_2\{Ph_2PCH_2S(O)Me\}\{MeOH\}]$ 185
G.H. Quek, P.-H. Leung, K.F. Mok (Singapore, Singapore)

Technical Note 189

Conference Calendar 190

Author Index of Volume 239 193

The publisher encourages the submission of articles in electronic form thus saving time and avoiding rekeying errors. A leaflet describing our requirements is available from the publisher upon request.